

Measure #126: Diabetic Foot and Ankle Care, Peripheral Neuropathy: Neurological Evaluation

DESCRIPTION:

Percentage of patients aged 18 years and older with a diagnosis of diabetes mellitus who had a neurological examination of their lower extremities

INSTRUCTIONS:

This measure is to be reported a minimum of once per reporting period for patients with diabetes mellitus seen during the reporting period. This measure may be reported by non-MD/DO clinicians who perform the quality actions described in the measure based on the services provided and the measure-specific denominator coding.

This measure is reported using G-codes:

ICD-9 diagnosis codes, CPT codes, and patient demographics (age, gender, etc.) are used to identify patients who are included in the measure's denominator. G-codes are used to report the numerator of the measure.

When reporting the measure, submit the appropriate denominator code(s) and the appropriate numerator G-code.

NUMERATOR:

Patients who had a lower extremity neurological exam performed at least once within 12 months

Definition: A lower extremity neurological exam consists of a documented evaluation of motor and sensory abilities including reflexes, vibratory, proprioception, sharp/dull and 5.07 filament detection.

Numerator Coding:

Lower Extremity Neurological Exam Performed

G8404: Lower extremity neurological exam performed and documented

OR

Lower Extremity Neurological Exam not Performed for Documented Reasons

G8406: Clinician documented that patient was not an eligible candidate for lower extremity neurological exam measure

OR

Lower Extremity Neurological Exam not Performed

G8405: Lower extremity neurological exam not performed

DENOMINATOR:

All patients aged 18 years and older with a diagnosis of diabetes mellitus

Denominator Coding:

An ICD-9 diagnosis code for diabetes mellitus and a CPT code are required to identify patients for denominator inclusion.

ICD-9 diagnosis codes: 250.00, 250.01, 250.02, 250.03, 250.10, 250.11, 250.12, 250.13, 250.20, 250.21, 250.22, 250.23, 250.30, 250.31, 250.32, 250.33, 250.40, 250.41, 250.42, 250.43, 250.50, 250.51, 250.52, 250.53, 250.60, 250.61, 250.62, 250.63, 250.70, 250.71, 250.72, 250.73, 250.80, 250.81, 250.82, 250.83, 250.90, 250.91, 250.92, 250.93

AND

CPT codes: 10060, 10061, 10180, 11000, 11040, 11041, 11042, 11043, 11044, 11055, 11056, 11057, 11719, 11720, 11721, 11730, 11740, 97001, 97002, 99201, 99202, 99203, 99204, 99205, 99211, 99212, 99213, 99214, 99215, 99304, 99305, 99306, 99307, 99308, 99309, 99310, 99324, 99325, 99326, 99327, 99328, 99334, 99335, 99336, 99337, 99341, 99342, 99343, 99344, 99345, 99347, 99348, 99349, 99350

RATIONALE:

Foot ulceration is the most common single precursor to lower extremity amputations among persons with diabetes. Treatment of infected foot wounds accounts for up to one-quarter of all inpatient hospital admissions for people with diabetes in the United States. Peripheral sensory neuropathy in the absence of perceived trauma is the primary factor leading to diabetic foot ulcerations. Approximately 45-60% of all diabetic ulcerations are purely neuropathic. Other forms of neuropathy may also play a role in foot ulcerations. Motor neuropathy resulting in anterior crural muscle atrophy or intrinsic muscle wasting can lead to foot deformities such as foot drop, equinus, and hammertoes. In people with diabetes, 22.8% have foot problems -- such as amputations and numbness -- compared with 10% of nondiabetics. Over the age of 40 years old, 30% of people with diabetes have loss of sensation in their feet.

CLINICAL RECOMMENDATION STATEMENTS:

Recognizing important risk factors and making a logical, treatment-oriented assessment of the diabetic foot requires a consistent and thorough diagnostic approach using a common language. Without such a method, the practitioner is more likely to overlook vital information and to pay inordinate attention to less critical points in the evaluation. A useful examination will involve identification of key risk factors and assignment into appropriate risk category. Only then can an effective treatment plan be designed and implemented. (ACFAS/ACFAOM Clinical Practice Guidelines)