

Major Depressive Disorder (MDD)

Patients who have Major Depression Disorder who who meet DSM IV Criteria

PQRI Data Collection Sheet

Patient's Name	Practice Medical Record Number (MRN)	Birth Date (mm/dd/yyyy) / /	Gender <input type="checkbox"/> Male <input type="checkbox"/> Female
National Provider Identifier (NPI)		Date of Service	

Clinical Information

Billing Information

Step 1 Is patient eligible for this measure?			Code Required on Claim Form
	Yes	No	
Patient is aged 18 years and older.	<input type="checkbox"/>	<input type="checkbox"/>	Verify date of birth on claim form.
Patient has a new diagnosis or recurrent episode of major depressive disorder.	<input type="checkbox"/>	<input type="checkbox"/>	Refer to coding specifications document for list of applicable codes.
There is a CPT E/M Service Code for this visit.	<input type="checkbox"/>	<input type="checkbox"/>	
If No is checked for any of the above, STOP. Do not report a CPT category II code or a G-code.			
Step 2 Does patient also have the other requirements for this measure?			Code to be Reported on Line 24D of Paper Claim Form (or Service Line 24 of Electronic Claim Form)
	Yes	No	
Is patient undergoing active treatment for a new diagnosis or recurrent episode of Major Depressive Disorder?	<input type="checkbox"/>	<input type="checkbox"/>	If No (ie, patient is in remission), report only G8466 and STOP. If Yes , report G8467 and proceed to Step 3.
Step 3 Does patient meet the measure?			Code to be Reported on Line 24D of Paper Claim Form, if Yes (or Service Line 24 of Electronic Claim Form)
DSM IV™ Criteria ¹ for Major Depressive Disorder	Yes	No	
Documented	<input type="checkbox"/>	<input type="checkbox"/>	1040F
			If No is checked for the above, report 1040F-8P (DSM IV™ Criteria for major depressive disorder not documented at the initial evaluation, reason not otherwise specified.)

¹DSM-IV™ criteria includes presence of depressed mood, marked diminished interest/pleasure, significant weight loss or weight gain, insomnia or hypersomnia, psychomotor agitation or retardation, fatigue or loss of energy, feelings of worthlessness, diminished ability to concentrate and recurrent suicidal ideation.