

Myelodysplastic Syndrome

Documentation of Iron Stores in Patients Receiving Erythropoietin Therapy

PQRI Data Collection Sheet

Patient's Name	Practice Medical Record Number (MRN)	Birth Date (mm/dd/yyyy) / /	Gender <input type="checkbox"/> Male <input type="checkbox"/> Female
National Provider Identifier (NPI)		Date of Service	

Clinical Information			Billing Information
Step 1 Is patient eligible for this measure?			
	Yes	No	Code Required on Claim Form
Patient is aged 18 years and older on date of encounter.	<input type="checkbox"/>	<input type="checkbox"/>	Verify date of birth on claim form.
Patient has a line item diagnosis of myelodysplastic syndrome.	<input type="checkbox"/>	<input type="checkbox"/>	Refer to coding specifications document for list of applicable codes. Codes determining a patient's eligibility must be reported on the same claim as the quality code(s) identified below.
There is a CPT E/M Service Code for this visit.	<input type="checkbox"/>	<input type="checkbox"/>	
If No is checked for any of the above, STOP. Do not report a CPT category II code.			
Step 2 Does patient also have the other requirements for this measure?			
	Yes	No	Code to be Reported on Line 24D of Paper Claim Form (or Service Line 24 of Electronic Claim Form)
Is patient receiving erythropoietin therapy ¹ ?	<input type="checkbox"/>	<input type="checkbox"/>	If No , report only 4095F and STOP. If Yes , report 4090F and proceed to Step 3.
Step 3 Does patient meet or have an acceptable reason for not meeting the measure?			
Iron Stores Prior to Initiating Erythropoietin Therapy	Yes	No	Code to be Reported on Line 24D of Paper Claim Form, if Yes (or Service Line 24 of Electronic Claim Form)
Documented ²	<input type="checkbox"/>	<input type="checkbox"/>	3160F
Not documented for the following reason: • System (eg, resources not available, other reason attributable to health care delivery system)	<input type="checkbox"/>	<input type="checkbox"/>	3160F-3P
Document reason here and in medical chart. _____ _____			If No is checked for all of the above, report 3160F-8P (Iron stores prior to initiating erythropoietin therapy not documented, reason not otherwise specified.)

¹For the purpose of this measure, erythropoietin therapy includes the following medications: epoetin and darbepoetin.

²Documentation of iron stores includes either: Bone marrow examination including iron stain OR serum iron measurement by ferritin or serum iron and TIBC.