

Rheumatoid Arthritis (RA) Measures Group

PQRI Data Collection Sheet*

Male Female

Patient's Name	Practice Medical Record Number (MRN)	Birth Date (mm/dd/yyyy)
National Provider Identifier (NPI)		Date of Encounter

Step 1 Preliminary reporting requirements

You must identify your intent to report the RA Measures Group by submitting the G-code specific to this measures group on the first patient claim (G8490: I intend to report the RA Measures Group). You do not need to resubmit the measures group-specific G-code on more than one claim.

Step 2 Determine patient eligibility

(Codes determining a patient's eligibility must be reported on the **same claim** as the quality code(s) identified in Step 3 below.)

	Yes	No	
Patient is aged 18 years and older on date of encounter.	<input type="checkbox"/>	<input type="checkbox"/>	Refer to date of birth listed above or on claim form.
Patient has a line item diagnosis of RA.	<input type="checkbox"/>	<input type="checkbox"/>	714.0, 714.1, 714.2, 714.81
There is a CPT E/M Service Code for an office visit/consultation, home visit, or for work related/medical disability evaluation.	<input type="checkbox"/>	<input type="checkbox"/>	99201, 99202, 99203, 99204, 99205, 99212, 99213, 99214, 99215, 99241, 99242, 99243, 99244, 99245, 99341, 99342, 99343, 99344, 99345, 99347, 99348, 99349, 99350, 99455, 99456

If **No** is checked for any of the above, STOP. This patient is not eligible for reporting on this measures group. Do not report a CPT category II code or G-code.

Step 3 Complete individual measures

Assessment and Classification of Disease Activity	Report one code for assessment/classification of disease activity OR one code for NOT assessed.	
PQRI Measure #177 <ul style="list-style-type: none"> reporting frequency: disease activity must be assessed, classified and reported once during the calendar year a standardized, systematic approach for evaluating the level of disease activity must be utilized. Standardized descriptive or numeric scales and/or composite indexes include but are not limited to: DAS28, SDAI, CDAI, RADA, RAPID 	Disease activity assessed and classified as low	<input type="checkbox"/> 3470F
	Disease activity assessed and classified as moderate	<input type="checkbox"/> 3471F
	Disease activity assessed and classified as high	<input type="checkbox"/> 3472F
	OR	
	Disease activity NOT assessed	<input type="checkbox"/> 3470F-8P
Assessment of Functional Status	Report one code for assessment of functional status OR one code for NOT assessed.	
PQRI Measure #178 <ul style="list-style-type: none"> reporting frequency: functional status must be assessed and reported once during the calendar year a standardized descriptive or numeric scale, standardized questionnaire, or notation of assessment of the impact of RA on patient activities of daily living must be utilized. Examples include but are not limited to: Health Assessment Questionnaire (HAQ), Modified HAQ, HAQ-2; American College of Rheumatology's Classification of Functional Status in RA 	Functional status assessed	<input type="checkbox"/> 1170F
	OR	
	Functional status NOT assessed	<input type="checkbox"/> 1170F-8P

continued on next page

*For additional information on the PQRI program and reporting on measures groups, please visit the CMS Web site at <http://www.cms.hhs.gov/pqri>.

Rheumatoid Arthritis (RA) Measures Group

continued from previous page

Assessment and Classification of Disease Prognosis	Report one code for assessment/classification of disease prognosis OR one code for NOT assessed.	
PQRI Measure #179 <ul style="list-style-type: none"> reporting frequency: disease prognosis must be assessed, classified and reported once during the calendar year classification should be based upon at a minimum the following: functional limitation, extraarticular disease (eg, vasculitis, Sjorgen's syndrome, RA lung disease, rheumatoid nodules), rheumatoid factor (RF) positivity, positive anti-cyclic citrullinated peptide (anti-CCP) antibodies (both characterized dichotomously, per CEP recommendation), and/or bony erosions by radiography RA patients with features of poor prognosis have active disease with high tender and swollen joint counts, often have evidence of radiographic erosions, elevated levels of RF and or anti-CCP antibodies, and an elevated erythrocyte sedimentation rate, and an elevated C-reactive protein level 	Disease prognosis assessed and classified as good	<input type="checkbox"/> 3476F
	Disease prognosis assessed and classified as poor	<input type="checkbox"/> 3475F
	OR	
	Disease prognosis NOT assessed	<input type="checkbox"/> 3475F-8P
Disease Modifying Anti-Rheumatic Drug (DMARD) Therapy	Report one code for DMARD therapy OR one code for NOT prescribed, dispensed, or administered.	
PQRI Measure #108 <ul style="list-style-type: none"> reporting frequency: DMARD therapy must be prescribed and reported once during the calendar year biologic DMARD therapy includes adalimumab, etanercept, infliximab, abatacept, anakinra and rituximab 	DMARD therapy prescribed, dispensed, or administered	<input type="checkbox"/> 4187F
	Not prescribed, dispensed, or administered for medical reasons [†] <ul style="list-style-type: none"> Document reason in medical chart 	<input type="checkbox"/> 4187F-1P
	OR	
	DMARD therapy NOT prescribed, dispensed, or administered	<input type="checkbox"/> 4187F-8P
Tuberculosis (TB) Screening Prior to First Course of Biologic DMARD Therapy	Report one code for biologic DMARD therapy status. If patient is receiving a first course of biologic DMARD therapy (excluding rituximab), you will also need to report one code for TB screening or one code for NOT performed.	
PQRI Measure #176 <ul style="list-style-type: none"> reporting frequency: TB screening must be performed and results interpreted 6 months prior to patient's first course of biologic DMARD therapy and reported once during the calendar year biologic DMARD therapy includes adalimumab, etanercept, infliximab, abatacept, anakinra (rituximab is excluded) patients are considered to be receiving a first course of therapy using a biologic DMARD only if they have never previously been prescribed or dispensed a biologic DMARD 	Patient not receiving a first course of biologic DMARD therapy (or biologic DMARD prescription is for rituximab)	<input type="checkbox"/> 4196F
	Patient receiving a first course of biologic DMARD therapy AND TB screening performed and results interpreted within 6 months prior to therapy	<input type="checkbox"/> 4195F AND 3455F
	Patient receiving a first course of biologic DMARD therapy AND TB screening not performed or results not interpreted for medical reasons (ie, patient positive for TB and documentation of past treatment; patient has recently completed a course of anti-TB therapy) <ul style="list-style-type: none"> Document reason in medical chart 	<input type="checkbox"/> 4195F and 3455F-1P
	OR	
	Patient receiving a first course of biologic DMARD therapy AND TB screening NOT performed or results NOT interpreted	<input type="checkbox"/> 4195F and 3455F-8P

continued on next page

[†]Medical reasons (eg, not indicated, contraindicated, other medical reason)

Rheumatoid Arthritis (RA) Measures Group

continued from previous page

Glucocorticoid Management		Report one code for glucocorticoid use (including dose) or ONE code for NOT assessed. If patient is receiving ≥ 10 mg daily prednisone (or equivalent) for longer than 6 months with improvement or no change in disease activity, you will also need to report one code for management plan OR one code for NOT documented.	
<p>PQRI Measure #180</p> <ul style="list-style-type: none"> reporting frequency: glucocorticoid use (and management plan, if appropriate) must be assessed and reported once during the calendar year 1 mg prednisone = 1 mg prednisolone; 5 mg cortisone; 4 mg hydrocortisone; 0.8 mg triamcinolone; 0.8 mg methylprednisolone; 0.15 mg dexamethasone; 0.15 mg betamethasone glucocorticoid management plan: attempt to taper steroids OR a new prescription for a non-glucocorticoid DMARD OR increase in dose of non-glucocorticoid DMARD dose for persistent RA disease activity at current or reduced dose 	<p>Assessment of glucocorticoid use (including dose) and management plan, if appropriate</p>	Patient not receiving glucocorticoid therapy	<input type="checkbox"/> 4192F
		Patient receiving < 10 mg daily prednisone (or equivalent) OR glucocorticoid use is for less than 6 months OR RA disease activity is worsening	<input type="checkbox"/> 4193F
		Patient receiving ≥ 10 mg daily prednisone (or equivalent) for longer than 6 months with improvement or no change in disease activity AND management plan documented	<input type="checkbox"/> 4194F AND 0540F
		Patient receiving ≥ 10 mg daily prednisone (or equivalent) for longer than 6 months with improvement or no change in disease activity AND management plan not documented for medical reasons (ie, glucocorticoid prescription is for a medical condition other than RA)	<input type="checkbox"/> 4194F AND 0540F-1P
		<p style="text-align: center;">OR</p> <p style="text-align: center;">(Report one of the following options)</p>	
Glucocorticoid use NOT assessed	<input type="checkbox"/> 4194F-8P		
Patient receiving ≥ 10 mg daily prednisone (or equivalent) for longer than 6 months with improvement or no change in disease activity AND management plan NOT documented	<input type="checkbox"/> 4194F AND 0540F-8P		

Step 4 Reporting Instructions

This measure can be reported for each eligible patient in one of two ways:

- Report the corresponding CPT category II code(s) as selected above for each of the six measures in the RA Measures Group

OR

- If **all** quality actions for the patient have been performed for each of the six measures in the RA Measures Group, **G8499** may be reported.

Note: G8499 is not appropriate for this patient if any CPT category II codes with the 8P modifier have been selected from Step 3.